



Effects of priming treatments on germination indices of soybean cultivars under osmotic stress

Masoud Radmanesh

Department of agriculture and Plant breeding, Islamic azad university, Zanjan, Iran

*Correspondence: Masoud_rad000000@yahoo.com Received: 01-05-2022, Revised: 18-06-2022, Accepted: 20-06-2022 e-Published: 23-06-2022

This study aimed to investigate the effects of priming treatments on germination indices of soybean cultivars under osmotic stress in a completely randomized design with a factorial arrangement of treatments in a botany laboratory located in Kordestan Province, Sanandaj, Iran. Experimental treatments included three treatments of priming (control and gibberellin at a concentration of 0.04%), cultivar (Clark and Hobbit), and osmotic stress (control and -1.5MPa) in three replications. The standard germination test was performed according to the guidelines presented by the International Seed Testing Association (ISTA) and some germination indices, including germination percentage, germination rate, germination uniformity, and mean germination time up to 10% germination (D10), were calculated at the end of the experiment using Germin program. In addition, some growth indices and enzymatic activities were also evaluated in this study. Results of analysis of variance indicated that the triple interaction of AxBxC had significant effects on all germination indices and the lowest coefficient of variation (equal to 7.55) was obtained for germination rate while its highest value (equal to 17.22) was calculated for germination uniformity index. Concerning the effects of priming treatment, it was observed that the highest values for indices of germination percentage and germination rate (with averages of 48% and 0.12, respectively) were obtained under the application of gibberellin. In other words, gibberellin treatment led to increases equal to 15.58 and 90.44% for germination percentage and germination rate indices compared to the control treatment, respectively. In addition, increasing osmotic stress levels had a significant inverse relationship with germination indices so that the application of -1.5 MPa osmotic stress significantly Decreased the germination percentage and germination rate of both Clark and Hobbit cultivars. In general, it was concluded that the application of gibberellin mitigated adverse effects of osmotic stress of -1.5 MPa and resulted in increasing percentage and rate of germination under stressful conditions.

Keywords: Germination, Gibberellin, Osmotic stress, Soybean

INTRODUCTION

Soybean (*Glycine max* L.), a self-pollinated diploid legume, is one of the most important oilseeds that makes up nearly 60% of the oilseed production in the world and its grains contain 40% protein, 20% oil, and high doses of B vitamins. The effects of climatic, environmental, and managerial factors on soybean, including moisture, soil temperature, air temperature, daylength, light intensity, soil pores, and soil conditions (soil fertility), are manifested through growth and developmental changes (Fataei, Monavari et al. 2010, Jia, Khayatnezhad et al. 2020, Arjaghi, Alasl et al. 2021, Bi, Chen et al. 2021, Chen, Khayatnezhad et al. 2021, Li, Mu et al. 2021). In general, soybeans are short-day (SD) and thermophilic plants that germinate at 8 to 10 ° C; however, the optimum temperature for its germination is around 30 ° C. In addition, it was reported that this plant is compatible with arid and semi-arid climate conditions, and its seeds mostly grow about 300 m elevation and daylength of 12 to 16

hours (Bi, Dan et al. 2021, Chen, Khayatnezhad et al. 2021, Cheng, Hong et al. 2021, Guo, She et al. 2021). Since soybean cultivars have relatively sensitive to day length, a 15-minute difference during the day may prevent flower growth and development in a unique cultivar (Tepperman and Dunsmuir 1990, Liu, Wang et al. 2021, Ma, Khayatnezhad et al. 2021, Peng, Khayatnezhad et al. 2021, Shi, Khayatnezhad et al. 2021). In addition to the above features, as a principle, early maturity groups of soybean cultivars need longer days for flowering, plant growth, and further development compared to late maturity groups (Jia, Khayatnezhad et al. 2020, Hou, Li et al. 2021, Huang, Wang et al. 2021).

Seed germination and emergence in seeds of many crops are the most sensitive stages in plants. In this regard, seed priming or priming technology, especially on soybean seeds, is one of the new methods of regulating seed germination in modern agriculture, in which seeds are first soaked in a solution for a definite time and dried

before emerging the radicle (Gholamin and Khayatnezhad 2020, Karasakal, Khayatnezhad et al. 2020, Ma, Ji et al. 2021, Zhu, Liu et al. 2021, Wang, Ma et al. 2022). Then, seed re-germination begins in the field at an appropriate time (as soon as conditions allow) using irrigation. Subsequently, the metabolic and physiological functions of the seeds continue, and plants begin to grow faster than expected conditions (Si, Gao et al. 2020, Garcia, Arif et al. 2021, Wang, Shang et al. 2021, Wang, Ye et al. 2021, Yin, Khayatnezhad et al. 2021).

On the other hand, since Iran is located in arid and semi-arid regions, water scarcity is one of the most important agricultural problems, and drought stress is a serious threat for global food security and plant production. According to the above definition of seed priming, the primary purpose of priming is to facilitate germination and promote the physiological and metabolic functions of seeds under particular environmental conditions, which leads to improved germination percentage and germination rate indices under challenging environments (Fataei, Varamesh et al. 2018, Gholamin and Khayatnezhad 2020, Sun and Khayatnezhad 2021). Also, priming minimizes the induced dormancy periods for various seeds by temperature and increases the temperature range required for seed germination (Abdi 2021, Alizadeh 2021, Karasakal 2021, Mohammadzadeh 2021, Radmanesh 2021). Overall, the term priming refers to the soaking of seeds in osmotic solutions with more negative water potential and ventilated, in which environmental conditions (e.g., temperature, light, and others) have significant effects on oxygen availability, the passage of time, seed drying, and control the seed-borne microbial contaminations during the dehydration process (Si, Gao et al. 2020, Guo, She et al. 2021, Khayatnezhad and Gholamin 2021, Li, Mu et al. 2021). In the meantime, gibberellic acid is a plant hormone and stimulant affecting the seed germination and dormancy in various plant species, which is employed as a priming solution for seeds. For more than two decades, seed priming has been applied as a standard method in many plants to alleviate the depressive effects of drought (Gholamin and Khayatnezhad 2020, Karasakal, Khayatnezhad et al. 2020). However, studies on the optimal reactions of primed and non-primed seeds are still under investigation, and seed priming continues until final evolution as a helpful technology. Accordingly, additional and further information becomes gradually available about primed seeds (Xiao, Yu et al. 2004, Zhao, Wang et al. 2021, Zheng, Zhao et al. 2021, Zhu, Saadati et al. 2021).

MATERIALS AND METHODS

This study aimed to investigate the effects of priming treatment on germination indices of soybean cultivars under osmotic stress. Therefore, an experiment was performed as a three-way factorial arrangement based on the completely randomized design in the fall season, 2020, in a botanical laboratory. Experimental treatments

included priming as factor A (a1: control and a2: gibberellin at a concentration of 0.04%), osmotic stress as factor B (b1: control and b2: -1.5MPa), and cultivar as factor C (c1: Clark and c2: Hobbit) in three replications. The standard germination tests were performed according to the ISTA Rules with slight modifications. Seed priming experiments (e.g., maximum germination, germination rate, and germination uniformity) were conducted as Petri experiments ((Hemmati, Fataei et al. 2019, Gholamin and Khayatnezhad 2020). Accordingly, three replicates of 25 seeds (5 series of 25 healthy seeds) were first counted to determine rate, percentage, and uniformity of germination. Next, seeds were disinfected with the carboxin-thiram fungicide to maintain and improve seed quality before planting in the culture medium. The seeds were then placed in Petri dishes (between two Whatman filters) and transferred to a growth chamber at 20 ± 1 °C. The number of germinated seeds scored three times per day for the time necessary to achieve the final number of germinated seeds. Hence, seeds were considered as germinated when their radicle length was more than or equal to 2 mm, and counting operations continued until the number of germinated seeds for each sample remained constant for three consecutive days. It should be noted that the Germin program, which calculates the indices of D10 (time up to 10% of maximum germination), D50 (time up to 50% of maximum germination), and D90 (time up to 90% of maximum germination), was employed for calculating traits of final germination percentage and germination rate (Mohammadi and Fataei 2019, Gholamin and Khayatnezhad 2021). This program calculates the above parameters for each replicate/seed treatment by interpolation from the progress of germination (%) versus the time curve. The germination rate, also called R50, was calculated in hours by the following equation (Soltani and Galeshi 2002).

Equation (1)

$$R50 = \frac{1}{D50}$$

Germination uniformity (GU) refers to the time required for 90% maximum germination subtracted by the time required for 10% maximum germination and is calculated by the following equation (Equation 2). In general, a lower value of this number indicates more uniform germination (simultaneous germination) of seeds.

Equation (2)

$$GU = D90 - D10$$

Statistical analysis

The obtained data were first normalized for different studied traits and then analyzed using SPSS 24 and MSTAT-C 21 statistical software. Also, the LSD test was used to compare means at $p < 0.05$, and charts were drawn using Excel software.

RESULTS AND DISCUSSION

Drought stress is one of the most important factors

limiting growth, development, and crop production in soybean and is considered a major challenging factor in many countries facing water scarcity (Nasehi and Fataei 2012, Yin, Khayatnezhad et al. 2021). Depending on the type of the plant and stress intensity, different methods have been used by numerous researchers and farmers to overcome water stress (Tao, Cui et al. 2021, Zhu, Saadati et al. 2021). Results of analysis of variance (Table 1) showed that factor A (priming), factor B (osmotic stress), and interactions of A × B, A × C, and A × B × C had significant effects on all germination indices. In addition, factor C (cultivar) and the interaction of B × C also caused significant differences in the obtained values of germination percentage, germination rate, and time up to 10% of maximum germination; however, both factor C and the interaction of B × C had no significant effects on germination uniformity. According to the data, the lowest coefficient of variation was (equal to 7.55) obtained for germination rate, while the highest value (with an average of 17.22) was observed for germination uniformity. In general, high values of coefficient of variation for GU and D10 indicate high effects of the environment on the above traits

In confirmation of the adverse effects of stressful conditions on germination indices, Yagmur and Kaydan(2008) revealed that osmotic stress reduced germination traits of triticale seeds. The negative impacts of osmotic stress on the initial growth of seeds of different plants were also investigated and confirmed by other researchers (Omran and Fataei 2018, Khayatnezhad and Gholamin 2020, Li, Mu et al. 2021, Xu, Ouyang et al. 2021). Xu et al. (2021) reported that drought stress and gibberellin priming significantly affected germination percentage, seedling length, dry weight of the treated seedlings, and seedling length and weight indices at $p < 0.01$. In addition, these traits were significantly affected by the interaction of gibberellin × drought stress ($p < 0.01$). Results presented by Khaliliaqdam and Mir-Mahmoodi (2013) also confirmed the significant effects of priming treatments on all germination indices of soybean seeds. In another study, Khalili et al. (2014) also reported that GP, R50, D10, D50, and D90 were significantly affected by temperature, water potential, and their interaction at $p < 0.01$.

Germination percentage

Seed germination depends on numerous environmental factors. In the meantime, water (and its availability) is one of the most influential factors in initiating seed germination and the physical and physiological processes of seed germination. Findings of some studies also indicated an increase in growth inhibitors, such as abscisic acid, and a decrease in growth-promoting hormones, e.g., auxins, gibberellins, and cytokinins, in plants under water scarcity conditions (Rajaei, Khalili-Arjaghi et al. 2020, Alizadeh 2021, Karasakal 2021, Mohammadzadeh 2021, Radmanesh 2021, Zhang,

Khayatnezhad et al. 2021). In the present study, the highest germination percentage (with an average of 90%) was recorded for the absence of gibberellin × absence of osmotic stress × Hobbit cultivar interaction, which placed in class A along with the absence of gibberellin × absence of osmotic stress × Clark cultivar interaction. In contrast, the germination percentage of seeds under treatments containing osmotic stress of -1.5 MPa was less than 10%. Results also showed that the germination percentage of seeds of Clark and Hobbit cultivars primed with gibberellin under -1.5 MPa conditions significantly increased compared to normal conditions. In other words, applying gibberellin reduced the adverse effects of osmotic stress (-1.5 MPa) and improved germination percentage (Fig. 1).




Figure1: Effects of tripartite interaction of priming × osmotic stress × cultivar (A×B×C) on germination percentage

Means sharing the same superscript are not significantly different from each other at $p < 0.05$ of the LSD test

Ghassemi-Golezani et al. (2008) reported an increase in the germination percentage of lentil seeds under priming with distilled water. They also stated that different responses of various genotypes to drought stress might depend on several factors, e.g., seed size, seed cover, and other characteristics. Decreasing water entrance into the seed under increasing drought stress can reduce hydraulic conductivity and thus decline the rate of physiological and metabolic processes involved in seed germination (Varamesh, Hosseini et al. 2014, Khayatnezhad and Gholamin 2020). Ghaffari et al. (2012) reported that although increasing stress levels resulted in a significant reduction in germination percentage, gibberellin treatment significantly increased seed germination percentage.

Table 1: Analysis of variance (mean squares) of the studied treatments on seed germination indices

Sources of variations	df	Germination percentage (G_{max})	Germination rate (R50)	Germination uniformity (GU)	time up to 10% of maximum germination (D10)
Rep	2	50.5 *	0.0000004 ^{ns}	111.2 ^{ns}	34.14 ^{ns}
Priming (Factor A)	1	82.08 **	0.00003 **	805.52 **	420.22 **
Osmotic stress (Factor B)	1	3842.44 **	0.0032 **	1132.8 **	1128.4 **
A×B interaction	1	1546.01 **	0.00005 **	1952.22 **	381.3 **
Cultivar (Factor C)	1	1208 **	0.0001 **	8.85 ^{ns}	2018.5 **
A×C interaction	1	752.7 **	0.00003 **	988.5 **	382.3 **
B×C interaction	1	985.2 **	0.00003 **	138.3 ^{ns}	2152.1 **
A×B×C interaction	1	914.4 **	0.000011 **	1528.22 **	346.54 **
E (Error)	51	19.521	0.0000003	68.32	25.22
Coefficient of variation (%)		8.88	7.55	17.22	16.34

ns: non-significant; * and **: significant at $p < 0.05$ and $p < 0.01$, respectively

Germination rate

Regulation and completion of metabolic stages involved in seed germination are the main reasons for increasing the germination rate of plant seeds under different priming treatments. In general, since non-primed seeds spend time performing phases I and II (rapid water imbibition by seed and reactivation of metabolism, respectively) longer, primed seeds are one step ahead of non-primed seeds in terms of germination and emergence stages (Garcia et al., 2021). Therefore, the cultivation of primed seeds, especially in unfavorable environmental conditions, leads to increased germination rate and germination percentage (Ren and Khayatnezhad 2021). Here, the highest germination rate was assigned to interactions of the absence of gibberellin×the absence of osmotic stress×Clark cultivar, the absence of gibberellin×the absence of osmotic stress×Hobbit cultivar, gibberellin treatment×the absence of osmotic stress×Clark cultivar, and gibberellin treatment×the absence of osmotic stress×Hobbit cultivar (Khayatnezhad and Gholamin 2021). Hence, all of the mentioned interactions were marked with the letter A and placed in the same group. According to the data, the application of gibberellin moderated adverse effects of drought stress-induced osmotic stress and led to increased germination rate in primed seeds of both Hobbit and Clark cultivars (Fig. 2). Decreased germination rate under stressful environmental conditions can also be directly correlated with reduced water absorption by seeds. As a whole, Hosseini and Rezvani Moghadam (2006) stated that seeds without priming solutions have low germination-related metabolic activities due to slow water uptake. Accordingly, the time required for radicle emergence and seed germination rate indices can be increased and decreased, respectively.




Figure 2: Effects of tripartite interaction of priming × osmotic stress × cultivar (A×B×C) on germination rate

Means sharing the same superscript are not significantly different from each other at $p < 0.05$ of the LSD test

Germination uniformity and time up to 10% of maximum germination (D10)

Germination uniformity was calculated based on the time required for 90% germination subtracted by the time required for 10% germination, namely D90-D10, in which scores with a lower value indicate a shorter time interval between 10% and 90% of seed germination and more favorable germination uniformity (Khayatnezhad and Nasehi 2021, Wang, Shang et al. 2021). Some researchers also stated that the lower absolute value for the obtained values indicated more uniformity for the germinated seeds (Sun, Lin et al. 2021, Li, Khayatnezhad et al. 2022, Wang, Khayatnezhad et al. 2022, Zhang, Khayatnezhad et al. 2022). Accordingly, since these seeds absorb water more rapidly under the same conditions, their germination rate and D10 occur higher and lower values, respectively. Based on the evidence, Rafiei (2001) stated that higher osmotic potential (less negative ψ_{π}) and less water absorption under stressful environmental conditions could lead to reducing germination rate, decreasing germination uniformity, and increasing time up to 10% of maximum germination; accordingly, seeds reached to final germination percentage in the longer growing periods. In this regard, our findings demonstrated that the lowest GU value

(namely the highest germination uniformity) was attributed to the interaction (combined treatment) of gibberellin \times osmotic stress \times Clark cultivar (Fig. 3).




Figure 3: Effects of tripartite interaction of priming \times osmotic stress \times cultivar (A \times B \times C) on germination uniformity

Means sharing the same superscript are not significantly different from each other at $p < 0.05$ of the LSD test




Figure 4: Effects of tripartite interaction of priming \times osmotic stress \times cultivar (A \times B \times C) on time up to 10% of maximum germination

Means sharing the same superscript are not significantly different from each other at $p < 0.05$ of the LSD test

CONCLUSION

In general, our findings showed that the highest germination percentage and germination rate (with average values of 48% and 0.12, respectively) were obtained for seeds primed with gibberellin and had increases of 18.58 and 90.44% for the above traits compared to the control treatment. On the other hand, osmotic stress had a significant inverse relationship with percentage and rate of germination indices in both studied cultivars. In other words, the application of -1.5 MPa osmotic stress caused notable reductions in germination percentage and germination rate indices compared to the normal conditions (absence of osmotic stress).

CONFLICT OF INTEREST

The authors declared that present study was performed in absence of any conflict of interest.

ACKNOWLEDGEMENT

This paper was from my own master thesis.

AUTHOR CONTRIBUTIONS

Masoud Radmanesh conducted, planned, Analyzed the data, wrote manuscript and interpreted the results and involved in manuscript preparation. All authors read and approved the final version.

Copyrights: © 2022@ author (s).

This is an open access article distributed under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

REFERENCES

- Abdi, M. M. (2021). "Evaluation of spring bread wheat lines/varieties in international observation nurseries and yield trials in moderat region of Iran." *Bioscience Research* 18(3): 2127-2134.
- Alizadeh, M. (2021). "Enhanced Imazethapyr Activity on Jimsonweed using natural Additives." *Bioscience Research* 18(4): 3080-3086.
- Alizadeh, M. (2021). "Protein profile and seeds storage proteins changes in wheat genotypes under control and drought stress conditions." *Bioscience Research* 18(3): 2164-2169.
- Arjaghi, S. K., M. K. Alasl, N. Sajjadi, E. Fataei and G. E. Rajaei (2021). "Green Synthesis of Iron Oxide Nanoparticles by RS Lichen Extract and its Application in Removing Heavy Metals of Lead and Cadmium." *Biological trace element research* 199(2).
- Bi, D., D. Chen, M. Khayatnezhad, Z. S. Hashjin, Z. Li and Y. Ma (2021). "molecular identification and genetic diversity in hypericum l.: a high value medicinal plant using rapd markers markers." *Genetika-Belgrade* 53(1): 393-405.
- Bi, D., C. Dan, M. Khayatnezhad, Z. sayyah hashjin, z. li and y. ma (2021). "molecular identification and genetic diversity in hypericum l.: a high value medicinal plant using rapd markers markers." *Genetika* 53(1): 393-405.
- Chen, W., M. Khayatnezhad and N. Sarhadi (2021). "Gene flow and population structure in Allochrusa (caryophylloideae, caryophyllaceae) with the use of molecular markers." *Genetika* 53(2): 799-812.
- Chen, W., M. Khayatnezhad and N. Sarhadi (2021). "protok gena i struktura populacije kod allochrusa (caryophylloideae, caryophyllaceae) pomocu molekularnih MARKERA." *Genetika-Belgrade* 53(2): 799-812.

- Cheng, X., X. Hong, M. Khayatnezhad and F. Ullah (2021). "Genetic diversity and comparative study of genomic DNA extraction protocols in *Tamarix* L. species." *Caryologia* 74(2): 131-139.
- Fataei, E., S. Monavari, A. Hasani, A. Karbasi and S. Mirbagheri (2010). "Heavy metal and agricultural toxics monitoring in Garasou river in Iran for water quality assessment." *Asian Journal of Chemistry* 22(4): 2991-3000.
- Fataei, E., S. Varamesh and S. Safavian (2018). "Effects of afforestation on carbon stocks in fandoghloo forest area." *Pakistan Journal of Agricultural Sciences* 55(3): 555-562.
- Garcia, D., S. Arif, Y. Zhao, S. Zhao, L. C. Ming and D. Huang (2021). "Seed Priming Technology as a Key Strategy to Increase Crop Plant Production under Adverse Environmental Conditions."
- GHASSEMI-GOLEZANI, K., A. Aliloo, M. Valizadeh and M. MOGHADDAM (2008). "Effects of hydro and osmo-priming on seed germination and field emergence of lentil (*Lens culinaris* Medik.)." *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 36(1): 29-33.
- Gholamin, R. and M. Khayatnezhad (2020). "Assessment of the Correlation between Chlorophyll Content and Drought Resistance in Corn Cultivars (*Zea Mays*)." *Helix* 10(5): 93-97.
- Gholamin, R. and M. Khayatnezhad (2020). "The effect of dry season stretch on Chlorophyll Content and RWC of Wheat Genotypes (*Triticum Durum* L.)." *Bioscience Biotechnology Research Communications* 13(4): 1829-1833.
- Gholamin, R. and M. Khayatnezhad (2020). "Study of Bread Wheat Genotype Physiological and Biochemical Responses to Drought Stress." *Helix* 10(5): 87-92.
- Gholamin, R. and M. Khayatnezhad (2020). "The Study of Path Analysis for Durum Wheat (*Triticum durum* Desf.) Yield Components." *Bioscience Biotechnology Research Communications* 13(4): 2139-2144.
- Gholamin, R. and M. Khayatnezhad (2021). "Impacts of PEG-6000-induced Drought Stress on Chlorophyll Content, Relative Water Content (RWC), and RNA Content of Peanut (*Arachis hypogaea* L.) Roots and Leaves." *Bioscience Research* 18(1): 393-402.
- Guo, L.-N., C. She, D.-B. Kong, S.-L. Yan, Y.-P. Xu, M. Khayatnezhad and F. Gholinia (2021). "Prediction of the effects of climate change on hydroelectric generation, electricity demand, and emissions of greenhouse gases under climatic scenarios and optimized ANN model." *Energy Reports* 7: 5431-5445.
- Hemmati, S., E. Fataei and A. A. Imani (2019). "Effects of Source Separation Education on Solid Waste Reduction in Developing Countries (A Case Study: Ardabil, Iran)." *The Journal of Solid Waste Technology and Management* 45(3): 267-272.
- HOSSEINI, H. and M. P. REZVANI (2006). "Effect of water and salinity stress in seed germination on Isabgol (*Plantago ovata*)."
- Hou, R., S. Li, M. Wu, G. Ren, W. Gao, M. Khayatnezhad and F. Gholinia (2021). "Assessing of impact climate parameters on the gap between hydropower supply and electricity demand by RCPs scenarios and optimized ANN by the improved Pathfinder (IPF) algorithm." *Energy* 237: 121621.
- Hu, W., R. Kaminski, F. Yang, Y. Zhang, L. Cosentino, F. Li, B. Luo, D. Alvarez-Carbonell, Y. Garcia-Mesa and J. Karn (2014). "RNA-directed gene editing specifically eradicates latent and prevents new HIV-1 infection." *Proceedings of the National Academy of Sciences* 111(31): 11461-11466.
- Huang, D., J. Wang and M. Khayatnezhad (2021). "Estimation of Actual Evapotranspiration Using Soil Moisture Balance and Remote Sensing." *Iranian Journal of Science and Technology, Transactions of Civil Engineering*: 1-8.
- Jia, Y., M. Khayatnezhad and S. Mehri (2020). "POPULATION DIFFERENTIATION AND GENE FLOW IN *Erodium cicutarium*: A POTENTIAL MEDICINAL PLANT." *Genetika-Belgrade* 52(3): 1127-1144.
- Jia, Y., M. Khayatnezhad and S. Mehri (2020). "Population differentiation and gene flow in *Erodium cicutarium*: A potential medicinal plant." *Genetika* 52(3): 1127-1144.
- Karasakal, A. (2021). "Composted organic manure replacement probability using chemical fertilizer in organic Safflower farming." *Bioscience Research* 18(4): 3115-3124.
- Karasakal, A. (2021). "Production of Wheat Seed Through Various Plantation Practices in Maternal Environment." *Bioscience Biotechnology Research Communications* 14(2): 549-552.
- Karasakal, A., M. Khayatnezhad and R. Gholamin (2020). "The Durum Wheat Gene Sequence Response Assessment of *Triticum durum* for Dehydration Situations Utilizing Different Indicators of Water Deficiency." *Bioscience Biotechnology Research Communications* 13(4): 2050-2057.
- Karasakal, A., M. Khayatnezhad and R. Gholamin (2020). "The Effect of Saline, Drought, and Presowing Salt Stress on Nitrate Reductase Activity in Varieties of *Eleusine coracana* (Gaertn)." *Bioscience Biotechnology Research Communications* 13(4): 2087-2091.
- Khayatnezhad, M. and R. Gholamin (2020). "A Modern Equation for Determining the Dry-spell Resistance of Crops to Identify Suitable Seeds for the Breeding Program Using Modified Stress Tolerance Index (MSTI)." *Bioscience Biotechnology Research Communications* 13(4): 2114-2117.
- Khayatnezhad, M. and R. Gholamin (2020). "Study of Durum Wheat Genotypes' Response to Drought

- Stress Conditions." *Helix* 10(5): 98-103.
- Khayatnezhad, M. and R. Gholamin (2021). "The Effect of Drought Stress on the Superoxide Dismutase and Chlorophyll Content in Durum Wheat Genotypes." *Advancements in Life Sciences* 8(2): 119-123.
- Khayatnezhad, M. and R. Gholamin (2021). "Impacts of Drought Stress on Corn Cultivars (*Zea mays* L.) At the Germination Stage." *Bioscience Research* 18(1): 409-414.
- Khayatnezhad, M. and F. Nasehi (2021). "Industrial Pesticides and a Methods Assessment for the Reduction of Associated Risks: A Review." *Advancements in Life Sciences* 8(2): 202-210.
- Li, A., X. Mu, X. Zhao, J. Xu, M. Khayatnezhad and R. Lalehzari (2021). "Developing the non-dimensional framework for water distribution formulation to evaluate sprinkler irrigation." *Irrigation and Drainage* 70(4): 659-667.
- Li, A., X. Mu, X. Zhao, J. Xu, M. Khayatnezhad and R. Lalehzari (2021). "Developing the non-dimensional framework for water distribution formulation to evaluate sprinkler irrigation." *Irrigation and Drainage*.
- Li, W., M. Khayatnezhad and A. Davarpanah (2022). "Statistical Analysis of Treated Flow-Back Water Measurements: An Industrial Insight for a Shale Reservoir." *Geofluids* 2022.
- Liu, S., Y. Wang, Y. Song, M. Khayatnezhad and A. A. Minaeifar (2021). "Genetic variations and interspecific relationships in *Salvia* (Lamiaceae) using SCoT molecular markers." *Caryologia* 74(3): 77-89.
- Ma, A., J. Ji and M. Khayatnezhad (2021). "Risk-constrained non-probabilistic scheduling of coordinated power-to-gas conversion facility and natural gas storage in power and gas based energy systems." *Sustainable Energy Grids & Networks* 26.
- Ma, S., M. Khayatnezhad and A. A. Minaeifar (2021). "Genetic diversity and relationships among *Hypericum* L. species by ISSR Markers: A high value medicinal plant from Northern of Iran." *Caryologia* 74(1): 97-107.
- Mir-Mahmoodi, T., N. Khaliliaqdam, A. Pirmani, S. Yazdan-Sta and S. Sharafi (2013). "Effects of hydro priming on laboratory indexes and field performance of sunflower seeds." *International Journal of AgriScience* 3(12): 904-911.
- Mohammadi, M. and E. Fataei (2019). "Comparative life cycle assessment of municipal wastewater treatment systems: lagoon and activated sludge." *Caspian Journal of Environmental Sciences* 17(4): 327-336.
- Mohammadzadeh, S. (2021). "The impact of trellising strategies on the production and quality of determinate tomato types." *Bioscience Research* 18(4): 3264-3271.
- Mohammadzadeh, S. (2021). "The presence of Nickel Sulphate in soil influences the synthesis of flavonoids and phenols in garlic (*Allium sativum* L.), changing its allelopathic potential." *Bioscience Research* 18(4): 3125-3131.
- Nasehi, F. and E. Fataei (2012). "Measurement of residue levels of agro-chemicals in water and sediment of Aras River." *Journal of Food, Agriculture & Environment* 10(1): 933-936.
- Omrani, M. and E. Fataei (2018). "Synthesizing Colloidal Zinc Oxide Nanoparticles for Effective Disinfection; Impact on the Inhibitory Growth of *Pseudomonas aeruginosa* on the Surface of an Infectious Unit." *Polish Journal of Environmental Studies* 27(4).
- Peng, X., M. Khayatnezhad and L. J. Ghezjeljheidan (2021). "RAPD PROFILING IN DETECTING GENETIC VARIATION IN *Stellaria* L. (Caryophyllaceae)." *Genetika-Belgrade* 53(1): 349-362.
- Radmanesh, M. (2021). "The Combined and Isolated Effect of Spinosad and Nuclear Polyhedrosis Virus on the Mediterranean Brocade *Spodoptera littoralis* in laboratory Conditions." *Bioscience Biotechnology Research Communications* 14(2): 692-696.
- Radmanesh, M. (2021). "Impacts of salinity, *Pseudomonas* and salicylic acid on seed bean physiological properties and yield (*Phaseolus vulgaris*)." *Bioscience Research* 18(4): 3023-3030.
- Rafiei, S.-R., H. A. Toliyat, R. Ghazi and T. Gopalathnam (2001). "An optimal and flexible control strategy for active filtering and power factor correction under non-sinusoidal line voltages." *IEEE Transactions on Power Delivery* 16(2): 297-305.
- Rajaei, G. E., S. Khalili-Arjaghi, E. Fataei, N. Sajjadi and M. Kashefi-Alasl (2020). "Fabrication and characterization of polymer-based nanocomposite membrane modified by magnetite nanoparticles for Cd $^{2+}$ and Pb $^{2+}$ removal from aqueous solutions." *Comptes Rendus. Chimie* 23(9-10): 563-574.
- Ren, J. and M. Khayatnezhad (2021). "Evaluating the stormwater management model to improve urban water allocation system in drought conditions." *Water Supply*.
- Shi, B., M. Khayatnezhad and A. Shakoor (2021). "The interacting effects of genetic variation in *Geranium* subg. *Geranium* (Geraniaceae) using scot molecular markers." *Caryologia* 74(3): 141-150.
- Si, X., L. Gao, Y. Song, M. Khayatnezhad and A. A. Minaeifar (2020). "Understanding population differentiation using geographical, morphological and genetic characterization in *Erodium cicutarium*." *Indian Journal of Genetics and Plant Breeding* 80(4): 459-+.
- Si, X., L. Gao, Y. Song, M. Khayatnezhad and A. A. Minaeifar (2020). "Understanding population differentiation using geographical, morphological and genetic characterization in *Erodium cicutarium*." *Indian J. Genet* 80(4): 459-467.
- Soltani, A. and S. Galeshi (2002). "Importance of rapid canopy closure for wheat production in a temperate

- sub-humid environment: experimentation and simulation." *Field Crops Research* 77(1): 17-30.
- Sun, Q., D. Lin, M. Khayatnezhad and M. Taghavi (2021). "Investigation of phosphoric acid fuel cell, linear Fresnel solar reflector and Organic Rankine Cycle polygeneration energy system in different climatic conditions." *Process Safety and Environmental Protection* 147: 993-1008.
- Sun, X. and M. Khayatnezhad (2021). "Fuzzy-probabilistic modeling the flood characteristics using bivariate frequency analysis and alpha-cut decomposition." *Water Supply* 21(8): 4391-4403.
- Tao, Z., Z. Cui, J. Yu and M. Khayatnezhad (2021). "Finite Difference Modelings of Groundwater Flow for Constructing Artificial Recharge Structures." *Iranian Journal of Science and Technology, Transactions of Civil Engineering*.
- Tepperman, J. M. and P. Dunsmuir (1990). "Transformed plants with elevated levels of chloroplastic SOD are not more resistant to superoxide toxicity." *Plant molecular biology* 14(4): 501-511.
- Varamesh, S., S. M. Hosseini, F. K. Behjou and E. Fataei (2014). "The impact of land afforestation on carbon stocks surrounding Tehran, Iran." *Journal of forestry research* 25(1): 135-141.
- Wang, C., Y. Shang and M. Khayatnezhad (2021). "Fuzzy Stress-based Modeling for Probabilistic Irrigation Planning Using Copula-NSPSO." *Water Resources Management* 35(14): 4943-4959.
- Wang, C., Y. Shang and M. Khayatnezhad (2021). "Fuzzy Stress-based Modeling for Probabilistic Irrigation Planning Using Copula-NSPSO." *Water Resources Management*.
- Wang, H., M. Khayatnezhad and N. Youssefi (2022). "Using an optimized soil and water assessment tool by deep belief networks to evaluate the impact of land use and climate change on water resources." *Concurrency and Computation-Practice & Experience* 34(10).
- Wang, J., Q. Ye, C. Wang, T. Zhang, X. Shi, M. Khayatnezhad and A. Shakoor (2021). "Palynological analysis of genus *Geranium* (Geraniaceae) and its systematic implications using scanning electron microscopy." *Caryologia* 74(3): 31-43.
- Wang, S., J. Ma, W. Li, M. Khayatnezhad and B. D. Rouyendegh (2022). "An optimal configuration for hybrid SOFC, gas turbine, and Proton Exchange Membrane Electrolyzer using a developed Aquila Optimizer." *International Journal of Hydrogen Energy* 47(14): 8943-8955.
- Xiao, B., Z. Yu, W. Huang, J. Song and P. a. Peng (2004). "Black carbon and kerogen in soils and sediments. 2. Their roles in equilibrium sorption of less-polar organic pollutants." *Environmental science & technology* 38(22): 5842-5852.
- Xu, Y.-P., P. Ouyang, S.-M. Xing, L.-Y. Qi, M. Khayatnezhad and H. Jafari (2021). "Optimal structure design of a PV/FC HRES using amended Water Strider Algorithm." *Energy Reports* 7: 2057-2067.
- Yagmur, M. and D. Kaydan (2008). "Alleviation of osmotic stress of water and salt in germination and seedling growth of triticale with seed priming treatments." *African Journal of Biotechnology* 7(13).
- Yin, J., M. Khayatnezhad and A. Shakoor (2021). "evaluation of genetic diversity in geranium (geraniaceae) using rapd marker." *Genetika* 53(1): 363-378.
- Yin, J., M. khayatnezhad and a. shakoor (2021). "evaluation of genetic diversity in geranium (geraniaceae) using rapd marker." *Genetika-Belgrade* 53(1): 363-378.
- Zhang, H., M. Khayatnezhad and A. Davarpanah (2021). "Experimental investigation on the application of carbon dioxide adsorption for a shale reservoir." *Energy Science & Engineering* n/a(n/a).
- Zhang, J., M. Khayatnezhad and N. Ghadimi (2022). "Optimal model evaluation of the proton-exchange membrane fuel cells based on deep learning and modified African Vulture Optimization Algorithm." *Energy Sources Part a-Recovery Utilization and Environmental Effects* 44(1): 287-305.
- Zhao, Y., H. Wang, W. Liang, M. Khayatnezhad and Faisal (2021). "GENETIC DIVERSITY AND RELATIONSHIPS AMONG *SALVIA* SPECIES BY ISSR MARKERS." *Genetika-Belgrade* 53(2): 559-574.
- Zheng, R., S. Zhao, M. Khayatnezhad and S. A. Shah (2021). "Comparative study and genetic diversity in *Salvia* (Lamiaceae) using RAPD Molecular Markers." *Caryologia* 74(2): 45-56.
- Zhu, K., L. Liu, S. Li, B. Li, M. Khayatnezhad and A. Shakoor (2021). "Morphological method and molecular marker determine genetic diversity and population structure in *Allochrusa*." *Caryologia* 74(2): 121-130.
- Zhu, P., H. Saadati and M. Khayatnezhad (2021). "Application of probability decision system and particle swarm optimization for improving soil moisture content." *Water Supply* 21(8): 4145-4152.